

B-Nine®/Florel® Spray in Propagation

- Use B-Nine® and Florel® PGRs together as a tank mix with a rate of 2500 ppm B-Nine / 500 ppm Florel (4 oz of B-Nine and 16 oz of Florel in 10 gallons of water).
- Apply as a spray in propagation, applying after root initiation and early growth (about 2 weeks after stick) depending on the crop. Florel can have a negative effect on initial rooting if used too soon.
- This early application can reduce the need for later applications in finished production, reducing or eliminating the delay in flowering that can occur with Florel.
- This combination spray may also eliminate the need for a pinch on some crops; trial pinch versus no pinch to see what works for you.
- Works well on Lantana, Impatiens, Calibrachoa, Petunia, Fuchsia, and Verbena.
- Do not use on New Guinea Impatiens, Torenia, Bacopa, or Euphorbia.



Dekko™ Star Rose Petunia
(New Jersey, Week 14, 2014)



Lanaj® Upright Blue with Eye Verbena
(New Jersey, Week 14, 2014)

Photos are either the property of Syngenta or used under agreement.

© 2018 Syngenta. **Important: Always read and follow label instructions. Some products may not be registered for sale or use in all states or countries. Please check with your state or local Extension Service to ensure registration status.** Some or all of the varieties listed herein may be protected under one or more of the following: Plant Variety Protection, United States Plant Patents, Utility Patents, and/or Plant Breeders' Rights and may not be propagated or reproduced without authorization. Trademarks used herein are trademarks of a Syngenta Group Company or their respective third party owners.